

Purpose

- Rapid identification of influenza virus infection is crucial during epidemics¹, allowing:
 - Hospital epidemiology interventions.
 - Rapid management of patient treatment (antivirals).
- However, classical surveillance models can only describe the status quo of an outbreak and do not allow predicting the future dynamics.
- The basic reproduction (R_0) number describes how many people are infected by a single infected individual e.g. $R_0 = 2$, indicates that 2 people are subsequently infected.

Datasets

University Hospital Basel, Influenza seasons 2013 – 2016


- All PCR confirmed cases (A+B) from the Hospital were included (**Figure 1**).
- PCR diagnostics was based on GeneXpert (Cepheid®).

Federal dataset of PCR-confirmed influenza cases reported in Switzerland

- All cases of Influenza have to be reported by law to the Federal Office of Health (**Figure 2**).
- Cases reported from 2007 to 2014 across age groups were available.
- We included all data from the cities Basel, Bern, and Zurich.

Methods

- A classic transmission-recovery model describes the spread.



- A statistical method was developed to estimate lambda and mu, and thus $R_0 = \lambda / \mu$.
- Assumption: Epidemics starts at week 0 with one case.
- In both datasets, the first six weeks were used for the R_0 calculation.

References

1 Miller MR, Peters TR et al J Infect Diseases 2015

Corresponding author

Adrian Egli, MD, PhD; Clinical Microbiology (adrian.egli@usb.ch)
University Hospital Basel, Petersgraben 4, 4031 Basel, Switzerland

Results




Figure 1. Absolute (A) and cumulative (B) cases of Influenza at the University Hospital Basel are shown. Based on the first six weeks, the R_0 was predicted using the computer algorithm.




Figure 2. Correlation of total cases per flu season and predicted R_0 value are shown. Data of the Federal Office of Health from three major Swiss cities were used.

Outlook of the project:

- Comparison of different algorithms to predict the basic reproduction number.
- Prospective validation of the algorithm over the next years.
- Comparison of R_0 between age groups.
- Implementation into epidemic management.

Summary and Conclusions

- The basic reproduction number (R_0) can be calculated 4-6 weeks after the first detection of an influenza case.
- The R_0 correlates with the cumulative amount of cases per season.
- In the near future such prediction algorithm based on real time influenza data can provide valuable information for epidemic management e.g. public health authorities.